



## Human IP-10 ELISA

Catalog Number EA-0505

(For Research Use Only)

### Introduction

IFN-gamma-inducible protein (IP-10) is a member of the chemokine family of cytokines and is induced in a variety of cells in response to interferon gamma and lipopolysaccharide. It is secreted by a number of cells including monocytes, endothelial cells and fibroblasts. IP-10 plays several roles, such as chemoattraction for monocytes/macrophages, T cells, NK cells, and dendritic cells, promotion of T cell adhesion to endothelial cells, antitumor activity, and inhibition of bone marrow colony formation. Several cell types in response to IFN- $\gamma$ . IP-10 acts as potent inhibitors of angiogenesis in vivo.

### Principle of the assay

IP-10 ELISA is based on the principle of a solid phase enzyme-linked immunosorbent assay. The assay utilizes rabbit anti-human IP-10 for immobilization on the microtiter wells and biotinylated rabbit anti-human IP-10 antibodies along with streptavidin conjugated to horseradish peroxidase (HRP) for detection. The test sample can react simultaneously with the two antibodies, resulting in the IP-10 molecules being sandwiched between the solid phase and enzyme-linked antibodies. After incubation, the wells are washed to remove unbound-labeled antibodies. An HRP substrate, TMB, is added to result in the development of a blue color. The color development is then stopped with the addition of Stop Solution changing the color to yellow. The concentration of IP-10 is directly proportional to the color intensity of the test sample. Absorbance is measured spectrophotometrically at 450 nm.



Diagram of ELISA

### Materials provided with the kit

Component	Qty	Store at
8x12 96-Well 12 strip Plate coated with rabbit anti-human IP-10 antibodies	1	4°C
Biotin labeled rabbit anti-human IP-10 antibodies	25 $\mu$ L	-20°C
Human IP-10 protein standard (200ng/ml)	10 $\mu$ L	-20°C
Streptavidin-HRP conjugate	50 $\mu$ L	4°C
1xDiluent buffer	40 mL	4°C
5X Assay wash buffer	40 mL	4°C
Substrate	10 mL	4°C
Stop solution	5 mL	4°C

### Material required but not provided


- Microplate reader capable of measuring absorbance at 450 nm
- Deionized or distilled water.

## Reagent preparation before starting experiment

- Dilute the 5x Assay wash buffer to 1x buffer  
40 ml 5x Assay wash buffer  
160 ml ddH<sub>2</sub>O
- Use serum-free conditioned media or original or 10-fold diluted sera. Sera can be diluted with 1X Diluent buffer. When serum-containing conditioned media is required, be sure to use serum as a control.
- Dilute 50 times of human IP-10 protein standard (200 ng/ml) with 1X Diluent buffer to 4000 pg/ml and then do a 2-fold serial dilutions by adding 4 ul of the diluted Human IP-10 protein standard in 200 ul 1X Diluent Buffer (See Step 2 below for detailed instruction).
- Dilute 400 times of biotin labeled rabbit anti-human IP-10 antibodies with 1X Diluent buffer before use.
- Dilute 200 times of streptavidin-HRP with 1X Diluent buffer before use.

## Assay procedure

1. Calculate the number of samples to decide how many strips need to be used.
2. Add 100  $\mu$ l of Standard, control, or sample per well and incubate for 1 hour at room temperature with gentle shaking.



- a. Add 200ul 1X Diluent buffer to the 1<sup>st</sup> well. Add 100ul 1X Diluent Buffer to the rest wells of strip.
- b. Add appropriate amount of protein recombinant (follow instruction in "Reagent Preparation")
- c. Mix dilutions in 1<sup>st</sup> well and transfer 100ul from the 1<sup>st</sup> well to the next dilution. (See picture) Incubate each well for 1 hr at room temperature with gentle shaking

3. Aspirate each well and wash by adding 200  $\mu$ l of 1X Assay wash buffer. Repeat the process three times for a total of three washes. Completely remove liquid at each wash. After the last wash, remove any remaining liquid by inverting the plate against clean paper towels.
4. Add 100  $\mu$ l of diluted biotin-labeled rabbit anti-human IP-10 antibodies to each well and incubate for 1 hour at room temperature with gentle shaking.
5. Repeat the aspiration/wash as in step 3.
6. Add 100  $\mu$ l of diluted streptavidin-HRP conjugate to each well and incubate for 45 min at room temperature with gentle shaking.

7. Repeat the aspiration/wash as in step 3.
8. Add 100  $\mu$ l of substrate to each well and incubate for 5-30 minutes.
9. Add 50  $\mu$ l of Stop solution to each well. The color in the wells should change from blue to yellow.
10. Determine the optical density of each well with a microplate reader at 450 nm within 30 minutes.

## Example of standard curve

